

IN THE CLAIMS

Please amend the claims as follows:

1-16 (Canceled)

17. (New) A method for use in a medical device, comprising:

providing a first plurality of sequential intervals based on cardiac electrical activity, with the plurality of atrial intervals including a plurality of shortest intervals, and a plurality of longest intervals, and a plurality of remaining intervals exclusive of the shortest and longest intervals;

determining a range statistic, a minimum interval statistic, and a dispersion statistic based on the plurality of remaining intervals, wherein the range statistic is based on the difference between a first and last occurring intervals of the remaining intervals, the minimum interval statistic is based on the shortest of the remaining intervals, and the dispersion statistic is based on a variance of the remaining intervals;

determining an assessment statistic based on the range statistic, the minimum interval statistic, and the dispersion statistic; and

selecting a cardiac therapy option based on the assessment statistic.

18. (New) The method of claim 17:

wherein the plurality of sequential intervals are atrial depolarization intervals; and

wherein selecting the a cardiac therapy option comprises:

determining whether the assessment statistic is greater than a threshold; and

selecting an atrial fibrillation therapy if the assessment statistic is determined to be greater than the threshold.

19. (New) The method of claim 17:

wherein the plurality of sequential intervals are atrial depolarization intervals; and

wherein selecting the cardiac therapy option comprises:

determining whether the assessment statistic is less than a threshold; and

selecting an atrial flutter therapy if the assessment statistic is determined to be less than the threshold.

20. (New) The method of claim 17, wherein selecting the cardiac therapy option comprises:
determining whether the assessment statistic is greater than a threshold; and
selecting a ventricular fibrillation therapy if the assessment statistic is determined to be
greater than the threshold.
21. (New) The method of claim 17, wherein selecting a cardiac therapy option comprises:
determining whether the assessment statistic is less than a threshold; and
selecting a ventricular tachycardia option if the assessment statistic is determined to be
less than the threshold.
22. (New) The method of claim 17, wherein the assessment statistic is inversely proportional
to the duration of the minimum interval and directly proportional to the range statistic and
to the dispersion statistic.
23. (New) The method of claim 17, wherein one or more of the intervals are composite
intervals.
24. (New) A machine-readable medium for controlling processing of a first plurality of
sequential intervals based on cardiac electrical activity, with the plurality of atrial
intervals including a plurality of shortest intervals, and a plurality of longest intervals, and
a plurality of remaining intervals exclusive of the shortest and longest intervals, the
medium comprising instructions for:
determining a range statistic, a minimum interval statistic, and a dispersion statistic based
on the plurality of remaining intervals, wherein the range statistic is based on the
difference between a first and last occurring intervals of the remaining intervals,
the minimum interval statistic is based on the shortest of the remaining intervals,
and the dispersion statistic is based on a variance of the remaining intervals;
determining an assessment statistic based on the range statistic, the minimum interval
statistic, and the dispersion statistic; and
selecting a cardiac therapy option based on the assessment statistic.

25. (New) The medium of claim 24:
wherein the plurality of sequential intervals are atrial depolarization intervals; and
wherein the instructions for selecting a cardiac therapy option comprise instructions for:
determining whether the assessment statistic is greater than a threshold; and
selecting an atrial fibrillation therapy if the assessment statistic is determined to be
greater than the threshold.
26. (New) The medium of claim 24:
wherein the plurality of sequential intervals are atrial depolarization intervals; and
wherein the instructions for selecting a cardiac therapy option comprise instructions for:
determining whether the assessment statistic is less than a threshold; and
selecting an atrial flutter therapy if the assessment statistic is determined to be less
than the threshold.
27. (New) The medium of claim 24, wherein the instructions for selecting the cardiac therapy
option comprise instructions for:
determining whether the assessment statistic is greater than a threshold; and
selecting a ventricular fibrillation therapy if the assessment statistic is determined to be
greater than the threshold.
28. (New) The medium of claim 24, wherein the instructions for selecting a cardiac therapy
option comprises:
determining whether the assessment statistic is less than a threshold; and
selecting a ventricular tachycardia option if the assessment statistic is determined to be
less than the threshold.
29. (New) The medium of claim 24, wherein the medium is an electronic memory.
30. (New) The medium of claim 24, wherein one or more of the intervals are composite
intervals.
31. (New) A method comprising:
calculating a set of two or more intervals based on data representative of an electrogram;
calculating a minimum interval based on the set of intervals;

calculating a range statistic based on a stability measurements for at least a subset of the intervals;

calculating a dispersion index based on a subset of the intervals;

calculating a number based on the minimum interval, the range statistic, the dispersion index; and

comparing the number to a threshold.

32. (New) The method of claim 31, wherein each stability measurement is a weighted average of differences between successive intervals.

33. (New) The method of claim 32, wherein the weighted average emphasizes relative importance of more recent intervals.

34. (New) The method of claim 31, wherein computing a number based on the minimum interval, the range statistic, and the dispersion index comprises computing the number based on an function having one of the following forms:

$$IDA_1 = K * \text{Range} * \text{Dispersion_index} * (\text{Min_interval})^{-1}$$

$$IDA_2 = K_1 * \text{Range} + K_2 * \text{Dispersion_index} + K_3 * (\text{Min_interval})^{-1},$$

where K, K₁, K₂, and K₃ denote constants; Range denotes the range statistic;

Dispersion_index denotes the dispersion index; and Min_interval denotes the minimum interval.

35. (New) The method of claim 31, wherein K and K₃ are each approximately 1.0, and wherein K₁ and K₂ are each approximately 0.0001.

Conclusion

Claims 1-16 have been canceled and claims 17-35 are added herein. Claims 17-35 are now pending. The Applicant respectfully requests that the supplemental preliminary amendment described herein be entered into the record prior to the examination and consideration of the above-identified application.

Respectfully Submitted,

MILTON M. MORRIS

By his Representatives,

SCHWEGMAN, LUNDBERG, WOESSNER & KLUTH, P.A.

P.O. Box 2938


Minneapolis, MN 55402

(612) 349-9593

Date

22 July 2004

By


Eduardo E. Drake

Reg. No. 40,594

CERTIFICATE UNDER 37 CFR § 1.8: The undersigned hereby certifies that this correspondence is being deposited with the United States Postal Service with sufficient postage as first class mail, in an envelope addressed to: Commissioner for Patents, MS: Missing Parts, P.O. Box 1450, Alexandria, VA 22313-1450, on this 22 day of July 2004.

Name

Lynne Mitchell-Becker

Signature

Lynne Mitchell-Becker